

# UCSF

## UC San Francisco Previously Published Works

### Title

Interpersonal prosodic correlation in frontotemporal dementia.

### Permalink

<https://escholarship.org/uc/item/5xz4n2rj>

### Journal

Annals of clinical and translational neurology, 6(7)

### ISSN

2328-9503

### Authors

Pressman, Peter S  
Ross, Elliott D  
Cohen, Kevin B  
et al.

### Publication Date

2019-07-01


### DOI

10.1002/acn3.50816

Peer reviewed

## BRIEF COMMUNICATION

# Interpersonal prosodic correlation in frontotemporal dementia

Peter S. Pressman<sup>1</sup> , Elliott D. Ross<sup>1</sup>, Kevin B. Cohen<sup>1</sup>, Kuan-Hua Chen<sup>2</sup>, Bruce L. Miller<sup>3</sup>, Lawrence E. Hunter<sup>1</sup>, Maria Luisa Gorno-Tempini<sup>3</sup> & Robert W. Levenson<sup>2</sup>

<sup>1</sup>Department of Neurology, Section of Behavioral Neurology and Neuropsychiatry, University of Colorado Denver, Anschutz Medical Campus, Academic Office Building 1, Mail Stop #B185, 12631 East 17th Avenue, Aurora, Colorado, 80045

<sup>2</sup>Berkeley Psychophysiology Lab, University of California, Berkeley, 4143 Tolman Hall, MC 5050, Berkeley, California, 94720-5050

<sup>3</sup>Memory and Aging Center, University of California, 675 Nelson Rising Ln, San Francisco, California, 94158

## Correspondence

Peter S. Pressman, Department of Neurology, Section of Behavioral Neurology and Neuropsychiatry, University of Colorado Denver, Anschutz Medical Campus, Academic Office Building 1, Mail Stop #B185, 12631 East 17th Avenue, Aurora, CO 80045. Tel: +1 303-724-0162; Fax: +1 303-724-2212; E-mail: peter.pressman@ucdenver.edu

## Funding information

Funding was provided through the American Brain Foundation and National Institutes of Health (NIH) grants P50AG023501, R01LM009254 and grant number P01AG019724.

Received: 5 March 2019; Revised: 8 May 2019; Accepted: 23 May 2019

*Annals of Clinical and Translational Neurology* 2019; 6(7): 1352–1357

doi: 10.1002/acn3.50816

## Abstract

Communication accommodation describes how individuals adjust their communicative style to that of their conversational partner. We predicted that interpersonal prosodic correlation related to pitch and timing would be decreased in behavioral variant frontotemporal dementia (bvFTD). We predicted that the interpersonal correlation in a timing measure and a pitch measure would be increased in right temporal FTD (rtFTD) due to sparing of the neural substrate for speech timing and pitch modulation but loss of social semantics. We found no significant effects in bvFTD, but conversations including rtFTD demonstrated higher interpersonal correlations in speech rate than healthy controls.

## Introduction

Communication accommodation theory describes how individuals adjust their communicative style to that of their partner in conversation. Communication accommodation can include linguistic (word choices), paralinguistic (pitch, tempo), and nonverbal (smiling, mutual gaze) features of communication. Accommodation establishes a paralinguistic conversational baseline, which may be broken to introduce new socio-emotional information into the conversation<sup>1</sup>. Lack of appropriate accommodation can have important social implications, as accommodation correlates with several measures of communication quality<sup>2</sup>.

Neurodegenerative disorders such as frontotemporal dementias (FTD) can cause deficits in communication and social interaction. Aberrant behavior often correlates with regions of neurodegeneration, thereby charting the neural underpinnings of social interactions. Assuming prosodic accommodation is a largely unconscious process, the phenomenon requires that the neural substrate for modulating the prosodic variable is intact, and that there is no recognized reason to break from the accommodative impulse. Paralinguistic pitch variation predominantly relies on a right hemispheric dorsal processing stream that ends in the orbitofrontal cortex. Timing depends on basal ganglia and cerebellar circuitry with some possible left predominance<sup>3</sup>. The neuroanatomy of communication

accommodation has been mapped onto the orbitofrontal cortex<sup>4</sup>, a region typically damaged in the behavioral variant of frontotemporal dementia (bvFTD). The right temporal variant of frontotemporal dementia relatively spares these areas, but has been associated with loss of social semantic knowledge and recognition of paralinguistic social signals<sup>5</sup>. Other types of FTD, such as the semantic and nonfluent variants of primary progressive aphasia (svPPA and nvPPA), primarily affect language while relatively sparing social and emotional functioning<sup>6</sup>.

We hypothesized that some types of FTD differ from healthy controls (i.e., conversational partners) in the extent to which prosody correlates between conversational partners. Specifically, we predicted that conversations including individuals with bvFTD would demonstrate less interpersonal correlation in pitch, intensity, and speech rate versus healthy controls due to a convergence of prosodic pathways in the orbitofrontal cortex. We predicted that correlation in speech rate and fundamental frequency modulation might be increased in the right temporal variant of frontotemporal dementia (rtFTD), because spared neural substrates for modulating speech rate and timing would permit unconscious accommodation, while loss of social semantics would diminish any recognized reason to break from that conversational baseline. For example, someone with rsvPPA would not recognize any need to slow or pause speech for emphasis, but would instead reflexively and inflexibly mimic his or her partner. We did not predict that interpersonal speech correlation

would be altered in Alzheimer's disease (AD), svPPA, or nvPPA.

## Methods

### Participants

About 74 patients with FTD (29 with bvFTD, 14 with rtFTD, 14 nvPPA, and 19 with svPPA), and 15 patients with Alzheimer's disease (AD) were compared to 170 healthy controls (patients' friends or family) in a recorded conversation lasting 10–15 min with a companion. Demographics and neuropsychological attributes for this population are listed in Table 1.

The Institutional Review Boards of the University of California, San Francisco, and the University of California, Berkeley, approved the study. All participants provided informed consent prior to participation. Prior to being assessed at the University of California, Berkeley, participants with a neurodegenerative illness underwent a detailed clinical evaluation, including a physical examination and neuropsychological testing at the University of California, San Francisco. Alzheimer's disease (AD) was established by National Institute on Aging – Alzheimer's Association criteria<sup>7</sup>, primary progressive aphasia (PPA), and bvFTD using consensus criteria<sup>6,8</sup>. Participants were diagnosed clinically with rtFTD based on clinical judgment and structural magnetic resonance imaging (MRI) data<sup>9,10</sup>.

**Table 1.** Participant demographics and neuropsychological characteristics

	HC	AD	bvFTD	nvPPA	rtFTD	svPPA
CDR	NA	1.1 ± 0.3	1.2 ± 0.6	0.6 ± 0.3*	0.9 ± 0.5	0.4 ± 0.3*
CDR Box	NA	6.1 ± 1.7	6.7 ± 2.9	2.6 ± 1.3*	5.5 ± 2.8	2.3 ± 1.7*
Education (years)	18.2 ± 1.3	15.9 ± 3.1	16.1 ± 2.9	17.4 ± 2.3	16.6 ± 2.3	17.4 ± 2.5
Age (years)	60.4 ± 10.8	60.8 ± 8.0	60.9 ± 8.8	67.4 ± 11.3	65.5 ± 5.7	63.7 ± 7.1
Handed (% right)	NA	86.7	93.1	92.3	92.3	94.7
Sex(%female)	61.7	40	34.5	50	42.9	21*
MMSE	NA	20.4 ± 5.2	25.1 ± 4.8	26.9 ± 3.2	25.6 ± 3.7	24.2 ± 3.9
BNT	NA	11.7 ± 4.3	11.6 ± 4.6	8.2 ± 2.7	8.0 ± 5.0	4.0 ± 2.9*
CVLT (30 sec)	NA	2.7 ± 2.9	4.8 ± 2.0	5.9 ± .6.6	4.0 ± 2.5	1.8 ± 2.3
CVLT (10 min)	NA	1.1 ± 2.5	4.1 ± 2.7	5.8 ± 2.8	2.7 ± 2.5	1.1 ± 1.9
Lexical fluency	NA	8.2 ± 6.0	5.9 ± 4.1	4.4 ± 3.3	6.1 ± 3.3	6.9 ± 3.8
Semantic fluency	NA	7.9 ± 4.7	9.6 ± 6.0	8.0 ± 6.5	9.5 ± 4.7	5.0 ± 3.0
Digits backwards	NA	2.5 ± 1.3	3.4 ± 1.5	3.6 ± 1.9	4.3 ± 1.7	4.4 ± 1.2
Benson copy	NA	9.4 ± 5.1	12.7 ± 5.0	10.4 ± 7.1	14.4 ± 2.9	15.2 ± 3.0
Benson recall (10 min)	NA	2.3 ± 2.8	6.4 ± 5.0	10.1 ± 3.6	5.7 ± 4.9	7.8 ± 4.2
Calculations	NA	2.9 ± 1.2	2.9 ± 1.7	3.5 ± 2.1	3.2 ± 2.4	4.4 ± 1.3
GDS	NA	8.0 ± 4.6	6.4 ± 6.7	4.4 ± 6.7	7.6 ± 9.8	8.2 ± 5.5

CD, clinical dementia rating scale; CDR box, CDR sum of box scores; MMSE, Mini Mental State Exam; BNT, Boston naming test abridged (15 items); CVLT, California Verbal Learning Test – II, GDS, geriatric depression scale.

\*Indicates  $P < 0.05$  difference.  $R^2 < 0.10$ .

## Task description

Procedures for obtaining samples of conversations were derived from those originally developed by Levenson and Gottman (1983)<sup>11</sup>. Couples were instructed to discuss a mutually selected area of continuing disagreement in their relationship for 10–15 min. Audio recordings of the conversations were obtained using unidirectional Shure lavalier microphones attached to each participant, recorded onto a single audio channel.

## Measures

Audio recordings were transformed into .wav files for further analysis. A spectral noise-gating algorithm was used to remove background noise<sup>12</sup>. Trained research assistants manually labeled all speech and nonspeech sounds for each speaker in Praat, an acoustic analysis program<sup>13</sup>. Environmental noises and nonspeech sounds were labeled for exclusion. Each labeled conversation was checked for quality before use. Using Praat, the following measures were extracted for each speaker: speech rate (syllables/second)<sup>14</sup>, coefficient of variation of fundamental frequency (i.e., pitch in Hz), and standard deviation of intensity (i.e., loudness in decibels).

## Statistical methods

A QQ plot was used to visually inspect data distributions of each measure. A cubic transformation was applied to achieve normative distributions for speech rate.

To compare interpersonal prosodic correlation slopes between diagnostic groups, regression between patient and partner speech with an interaction term for patient diagnosis was performed, adjusting for patient age and sex. Plots of residuals versus fitted values were inspected for heteroscedasticity. Regression was then repeated with robust standard errors as needed. All regressions with statistically significant results were assessed for influential

outliers. If potential outliers were identified, the regression was repeated without those outliers included. A *P* value of 0.05 and *R*<sup>2</sup> of 0.10 was set as being statistically and behaviorally significant, respectively.

## Results

Means and standard deviations for prosodic measurements in each group are listed in Table 2. An ANOVA revealed no differences between any diagnostic group in how either speakers or partners varied fundamental frequency or intensity (all *P* > 0.15). Differences were found in speech rate, but these did not meet the effect size criterion.

When comparing the extent to which patient's speech traits correlated with that of their partners across each group (i.e., the interaction term in the regression), conversations between patients with rtFTD and their partners had a markedly higher degree of positive speech rate correlation ( $\beta = 0.8$ , *P* = 0.004, 95%CI [0.26; 1.39], *R*<sup>2</sup> = 0.35, Fig. 1). Initial analysis suggested possible diminished interpersonal correlation of fundamental frequency variation in conversations including bvFTD ( $\beta = -0.45$ , *P* = 0.014, 95%CI [-0.8, -0.1], *R*<sup>2</sup> = 0.10). Further regression diagnostics suggested substantial heteroscedasticity in the residuals, as well as the possibility of results being driven by an extreme case. No differences in the interpersonal prosodic slopes were found between groups when using a more robust regression or after removing the potential outlier. We also did not find differing interpersonal prosodic correlations in conversations including nvPPA or svPPA.

## Discussion

We predicted that compared with healthy controls, the extent to which prosodic measures correlated between speakers would be lower for fundamental frequency, intensity, and rate in bvFTD, and increased for rate and


**Table 2.** Acoustic speech measures

	HC	AD	bvFTD	nvPPA	rtFTD	svPPA
CoVar of <i>F</i> <sub>0</sub>	20.0 ± 6.7	19.8 ± 4.5	21.3 ± 11.0	20.4 ± 7.7	18.6 ± 6.9	18.6 ± 6.3
Partner CoVar of <i>F</i> <sub>0</sub>	20.6 ± 10.1	19.0 ± 7.1	22.1 ± 7.7	18.4 ± 6.3	16.7 ± 4.1	18.6 ± 6.3
Speech rate (syl/sec)	3.3 ± 0.7	3.2 ± 0.5	3.2 ± 0.9*	2.75 ± 0.8*	3.7 ± 0.8	3.3 ± 0.6
Partner speech rate	3.3 ± 0.7	3.1 ± 0.5	3.4 ± 0.9	3.4 ± 0.7	3.8 ± 0.4	3.7 ± 0.5
Speech rate (cubed)	48.3 ± 21.6	34.3 ± 14.3	38.2 ± 20.8*	25.8 ± 19.3*	55.2 ± 31.0	38.3 ± 23.5
Partner speech rate (cubed)	39.4 ± 21.7	42.8 ± 16.4	47.3 ± 27.2	42.3 ± 22.3	58.9 ± 18.9*	54.0 ± 20.0*
Intensity SD (dB)	14.1 ± 2.6	15.1 ± 2.3	14.2 ± 2.9	13.8 ± 2.5	14.1 ± 2.4	14.9 ± 2.2
Partner intensity SD	14.4 ± 2.6	14.7 ± 1.4	14.0 ± 3.3	13.3 ± 3.4	13.7 ± 2.0	14.1 ± 2.1

CoVar of *F*<sub>0</sub>, coefficient of variation of fundamental frequency in Hertz. Speech rate is in syllables per second. Intensity as measured in decibels.

\*Values in which *P* values < 0.05.

## Speech rate compared to partner's speech rate



**Figure 1.** Correlation between participant and partner speech rate as measured in syllables per second of speech cubed. Although most forms of FTD as well as healthy controls show a consistent slope, conversations including rtFTD are associated with a significantly increased slope of correlation between speakers.

fundamental frequency in rtFTD. Contrary to what might be predicted based on reports of orbitofrontal involvement in accommodation, we found no robust evidence to support diminished prosodic correlation in bvFTD. As hypothesized, the slope of interpersonal correlation of speech rate was greater in conversations with patients with rtFTD than healthy controls.

Right frontotemporal dementia is thought to be associated with a loss of semantic knowledge for socially relevant concepts, such as reduced knowledge of social appropriateness, deficits in recognition of facial identity, and expression and reduced empathy<sup>5</sup>. Nevertheless, neural substrates for prosodic expression, particularly for timing, are relatively spared in rtFTD<sup>3</sup>. Therefore, our findings of increased interpersonal prosodic correlation for speech rate in conversations involving rtFTD likely relates to reduced recognition of the need to break from a reflexive imitation of the partner's prosodic style, with spared ability to imitate that partner's timing.

As expected, we did not find altered interpersonal prosodic correlation in svPPA or nfvPPA. This result may

seem surprising in nfvPPA, which is partially defined by nonfluency. The sustained ability of nonfluent patients with left frontal damage to mimic fluent speech stimuli, however<sup>15</sup>, is sufficiently well established to be the basis of therapeutic trials in nfvPPA<sup>16</sup>. In addition, contrary to some prior research on bvFTD<sup>17</sup>, we found no diagnostic group differences in standard deviation of fundamental frequency. This negative finding aligns with other research, however<sup>18</sup>, and may represent differences in statistical technique, the speech elicitation task, and/or in the sample selection (e.g., bvFTD is a heterogeneous disorder, and may be prone to unrecognized differences in prosodic accommodation between subtypes)<sup>19</sup>.

Our analytic approach was limited to the level of the entire conversation – we did not specify whether patients adjust their speech more to caregivers or vice versa. A different statistical approach is necessary to distinguish the extent to which each individual matches their speech to that of their partner over the course of conversations. This more detailed approach would provide individual (rather than conversational) correlation values, thereby

permitting correlations with other neuropsychological tests and neuroimaging studies. Further limitations of this study include the use of patients' friends and family members as healthy controls, with limited information available about that population to ensure generalizability. The successful prediction of an aspect of interpersonal prosodic correlation nevertheless suggests that speech accommodation may be a quantifiable behavioral marker of neurodegenerative disease.

## Acknowledgments

We thank the participants and staff at UCSF's Memory and Aging Center, the University of California, Berkeley's Psychophysiology Laboratory, the Tau Consortium, and the Consortium for Frontotemporal Dementia Research. We thank Dr. Celine De Looze for advising this research. Funding was provided through the American Brain Foundation and National Institutes of Health (NIH) grants P50AG023501, R01LM 009254 and grant number P01AG019724.

## Conflict of Interest

Peter S. Pressman reports no disclosures.

Kuan-Hua Chen reports no disclosures.

Elliott Ross reports no disclosures.

Bruce L. Miller has served as director on the medical advisory board of the Larry H. Hillblom Foundation, medical director of the John Douglas French Foundation, and scientific director of the Tau Consortium.

Elliott Ross reports no disclosures.

Kevin B. Cohen reports no disclosures.

Lawrence Hunter reports serves as a member of the Scientific Advisory Board of Medibio, Inc. and is a consultant to Somalogic, Inc.

Maria Luisa Gorno-Tempini reports no disclosures.

Robert W. Levenson reports no disclosures.

## Author Contributions

Peter S. Pressman – Study concept and design, analysis, and interpretation.

Kevin B. Cohen – Critical revision of the manuscript for important intellectual content.

Elliott Ross – Analysis and interpretation, critical revision of the manuscript for important intellectual content.

Bruce L. Miller – Study supervision, critical revision of the manuscript for important intellectual content.

Kuan-Hua Chen – Study concept and analysis, critical revision of the manuscript for important intellectual content.

Lawrence E. Hunter – Critical revision of the manuscript for important intellectual content.

Maria Luisa Gorno-Tempini – Study supervision, critical revision of the manuscript for important intellectual content.

Robert W. Levenson – Study supervision, critical revision of the manuscript for important intellectual content.

## References

1. Giles H, Coupland N, Coupland J. Accommodation theory: communication, context, and consequence. Contexts of accommodation: developments in applied sociolinguistics. Cambridge, England: Cambridge University Press, 1991:1–68.
2. Gregory S, Webster S, Huang G. Voice pitch and amplitude convergence as a metric of quality in dyadic interviews. *Lang Commun* 1993;13:195–217.
3. Fruhholz S, Trost W, Kotz SA. The sound of emotions—Towards a unifying neural network perspective of affective sound processing. *Neurosci Biobehav Rev* 2016;68:96–110.
4. Donohue WA, Sherry JL, Idzik P. Interaction dynamics predict successful negotiation in divorce mediation. *J Lang Soc Psychol* 2016;35:374–93.
5. Kamminga J, Kumfor F, Burrell JR, et al. Differentiating between right-lateralised semantic dementia and behavioural-variant frontotemporal dementia: an examination of clinical characteristics and emotion processing. *J Neurol Neurosurg Psychiatry* 2015;86:1082–1088.
6. Gorno-Tempini ML, Hillis AE, Weintraub S, et al. Classification of primary progressive aphasia and its variants. *Neurology* 2011;76:1006–14.
7. McKhann GM, Knopman DS, Chertkow H, et al. The diagnosis of dementia due to Alzheimer's disease: recommendations from the National Institute on Aging-Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. *Alzheimer's Dement* 2011;7:263–9.
8. Rasovsky K, Hodges JR, Knopman D, et al. Sensitivity of revised diagnostic criteria for the behavioural variant of frontotemporal dementia. *Brain* 2011;134(Pt 9):2456–77.
9. Josephs KA, Whitwell JL, Knopman DS, et al. Two distinct subtypes of right temporal variant frontotemporal dementia. *Neurology* 2009;73:1443–50.
10. Binney RJ, Henry ML, Babiak M, et al. Reading words and other people: a comparison of exception word, familiar face and affect processing in the left and right temporal variants of primary progressive aphasia. *Cortex* 2016;82:11.
11. Levenson RW, Gottman JM. Marital interaction: physiological linkage and affective exchange. *J Person Soc Psychol* 1983;45:587–97.
12. Team Audacity. How audacity noise reduction works. 2015 [updated March 30, 2014; cited 2015 May 1]; Available from: [http://wiki.audacityteam.org/wiki/How\\_Audacity\\_Noise\\_Reduction\\_Works](http://wiki.audacityteam.org/wiki/How_Audacity_Noise_Reduction_Works).

13. Boersma P, Weenink D. Praat 5.4.01 ed2014.
14. de Jong NH, Wempe T. Praat script to detect syllable nuclei and measure speech rate automatically. *Behav Res Methods* 2009;41:385–390.
15. Fridriksson J, Hubbard HI, Hudspeth SG, et al. Speech entrainment enables patients with Broca's aphasia to produce fluent speech. *Brain* 2012;135(Pt 12):3815–29.
16. Henry ML, Hubbard HI, Grasso SM, et al. Retraining speech production and fluency in non-fluent/agrammatic primary progressive aphasia. *Brain* 2018;141:1799–814.
17. Nevler N, Ash S, Jester C, et al. Automatic measurement of prosody in behavioral variant FTD. *Neurology* 2017;89:650–656.
18. Vogel AP, Poole ML, Pemberton H, et al. Motor speech signature of behavioral variant frontotemporal dementia: refining the phenotype. *Neurology* 2017;89:837–844.
19. Ranasinghe KG, Rankin KP, Pressman PS, et al. Distinct subtypes of behavioral variant frontotemporal dementia based on patterns of network degeneration. *JAMA Neurol* 2016;73:1078–88.